

# Teaching Neuro*Images*: Isolated sphenoiditis: An uncommon cause of abducens nerve palsy

Apostolos Safouris, Andrey Bragin, Chiara Mabiglia, et al. Neurology 2013;81;e176 DOI 10.1212/01.wnl.0000436936.24787.82

This information is current as of December 2, 2013

The online version of this article, along with updated information and services, is located on the World Wide Web at: http://www.neurology.org/content/81/23/e176.full.html

*Neurology* <sup>®</sup> is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2013 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.





### Section Editor Mitchell S.V. Elkind, MD, MS

Apostolos Safouris, MD Andrey Bragin, MD Chiara Mabiglia, MD Nikos Triantafyllou, MD Marie-Dominique Gazagnes, MD Georgios Tsivgoulis, MD, FESO

Correspondence to Dr. Safouris: safouris@yahoo.com

## Teaching Neuro*Images*: Isolated sphenoiditis

An uncommon cause of abducens nerve palsy

#### Figure Imaging



(A) Noncontrast brain CT was initially interpreted as normal but shows an unpneumatized left sphenoid sinus. (B) Coronal T2 brain MRI shows hyperintensity in the left sphenoid sinus, consistent with inflammation. (C) Axial fluid-attenuated inversion recovery MRI shows significant deviation of the nasal septum, a predisposing factor to sphenoiditis.

A 35-year-old woman presented with a headache lasting 2 weeks and complained of recent diplopia. Neurologic examination showed limited left eye abduction. Initial brain CT scan was considered normal (figure, A). Brain MRI showed left sphenoiditis (figure, B). The patient was treated with amoxicillin/clavulanic acid and corticosteroids with rapid symptom improvement which deferred endoscopy that could provide a bacteriologic diagnosis.

Isolated sphenoiditis represents fewer than 3% of sinusitis cases and rarely may be complicated by abducens nerve palsy due to the nerve's proximity to the sphenoid sinus at the level of the petrous apex and cavernous sinus.<sup>1</sup> Coronal T2 sequences on brain MRI may facilitate prompt diagnosis.<sup>2</sup>

#### AUTHOR CONTRIBUTIONS

Apostolos Safouris: study design, data collection, drafting and revising the manuscript. Andrey Bragin: data collection, critical comments during manuscript revision. Chiara Mabiglia: data collection, critical comments during manuscript revision. Nikos Triantafyllou: critical comments during manuscript revision. Marie-Dominique Gazagnes: data collection, critical comments during manuscript revision. Georgios Tsivgoulis: study design, drafting and revising the manuscript.

#### STUDY FUNDING

Dr. Georgios Tsivgoulis was supported by the European Regional Development Fund, Project FNUSA-ICRC (No. CZ.1.05/1.1.00/02.0123).

#### DISCLOSURE

The authors report no disclosures relevant to the manuscript. Go to Neurology.org for full disclosures.

#### REFERENCES

- Wang ZM, Kanoh N, Dai CF, et al. Isolated sphenoid sinus disease: an analysis of 122 cases. Ann Otol Rhinol Laryngol 2002;111:323–327.
- Sieskiewicz A, Lyson T, Olszewska E, Chlabicz M, Buonamassa S, Rogowski M. Isolated sphenoid sinus pathologies: the problem of delayed diagnosis. Med Sci Monit 2011;17:180–184.

Download teaching slides: www.neurology.org

> From the Stroke Unit, Department of Neurology (A.S., A.B., M.-D.G.), and the Department of Radiology (C.M.), Brugmann University Hospital, Brussels, Belgium; First Department of Neurology (N.T.), University of Athens, School of Medicine, "Eginition" University Hospital; Second Department of Neurology (G.T.), University of Athens, School of Medicine, "Attikon" University Hospital, Greece; and the International Clinical Research Center (G.T.), St. Anne's University Hospital, Brno, Czech Republic.

© 2013 American Academy of Neurology. Unauthorized reproduction of this article is prohibited.

## Teaching NeuroImages: Isolated sphenoiditis: An uncommon cause of abducens nerve palsy Apostolos Safouris, Andrey Bragin, Chiara Mabiglia, et al. Neurology 2013;81;e176 DOI 10.1212/01.wnl.0000436936.24787.82

Updated Information & Services	including high resolution figures, can be found at: http://www.neurology.org/content/81/23/e176.full.html
Supplementary Material	Supplementary material can be found at: http://www.neurology.org/content/suppl/2013/11/29/81.23.e176. DC1.html
References	This article cites 2 articles, 0 of which you can access for free at: http://www.neurology.org/content/81/23/e176.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): <b>All Headache</b> http://www.neurology.org//cgi/collection/all_headache <b>Diplopia (double vision)</b> http://www.neurology.org//cgi/collection/diplopia_double_vision
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://www.neurology.org/misc/addir.xhtml#reprintsus

## This information is current as of December 2, 2013

